

Heavy Quarks at Threshold: Recent Developments

Matthias Steinhauser

*II. Institut für Theoretische Physik, Universität Hamburg,
22761 Hamburg, Germany*

Recent developments in the treatment of a system of heavy quarks close to their production threshold are discussed. Particular emphasis is put on the next-to-next-to-next-to-leading order (N³LO) corrections to the energy levels and the wave function and their effect on the cross section $\sigma(e^+e^- \rightarrow t\bar{t})$. Furthermore, the resummation of logarithmic corrections within the framework of potential non-relativistic QCD (pNRQCD) are considered.

1 Introduction

One of the most important tasks of a future linear collider is the precise measurement of the cross section for the production of top quark pairs close to their production threshold. Next to the mass and the width of the top quark also the strong coupling and — in case the Higgs boson is not too heavy — also the top quark Yukawa coupling can be determined with a quite high accuracy. In order to match the expected experimental precision¹ it is important to compute higher-order quantum corrections to this process.

Due to the large mass and width top quarks do not form bound states. Instead, the resonance curve in the threshold region shows a local maximum below the nominal threshold which is essentially the remnant of a would-be $1S$ bound state.

The NNLO corrections to the cross section $e^+e^- \rightarrow t\bar{t}$ in the threshold region has been completed a few years ago by four independent groups. The comparison of the individual results is discussed in Ref.². The essential outcome of the combined effort is that no stability of the position of the peak is found in case the pole quark mass is used for the parametrization of the cross section. Thus a precise determination of the quark mass would not be possible. However, this changes if one switches to a short-distance mass which leads to a stabilization of the peak position and thus to a reliable extraction of the corresponding mass parameter. The normalization in the peak position, which is essentially determined by the wave function at the origin, still remains unstable after including NLO and NNLO corrections.

Shortly after the completion of the second order calculation there have been attempts to go beyond the fixed-order approximation and resum the logarithms in the velocity of the top quark. In Ref.³ an accuracy of 2-3% was claimed for the cross section of $t\bar{t}$ threshold production. However, subsequent

calculations of further next-to-next-to-leading logarithmic (NNLL) terms⁴, which had not been taken into account in Ref.³, casted serious doubts on this optimistic estimate. Thus, the full calculation of the NNLL corrections, which still remains elusive, is unavoidable to draw definite conclusions.

In these proceedings we want to report about phenomenological applications which are based on higher order results obtained within the framework of pNRQCD⁵. In particular, it was possible to evaluate the fixed-order N³LO corrections to the ground state energy^{6,7}. After the inclusion of the third-order corrections an impressive stabilization in the position of the peak is observed — even for on-shell quark masses. Thus this result can be used to determine the bottom quark mass from the $\Upsilon(1S)$ system and the top quark mass from the position of the peak in the cross section⁷. The latter will be discussed in Section 2. Furthermore, the quadratic⁸ and linear^{9,4} logarithms of the third-order corrections to the wave function have been determined. The inclusion of these terms seems to provide more stability in the normalization of the peak. However, for a definite conclusion, also the constant term at N³LO is needed.

In the meantime first results containing the summation of higher-order logarithms within pNRQCD became available for the spin-dependent part of the Hamiltonian. In particular the NLL corrections to the hyperfine splitting (HFS)^a have been evaluated in¹⁰. In Section 3 we will discuss how this result can be used to predict the mass of the η_b meson. In Ref.¹¹ the results of¹⁰ have been generalized to the non-equal mass case and applied to the HFS in the B_c system. For completeness we want to mention that recently the NNLL corrections to the ratio of the spin-1 and spin-0 mediated production and annihilation processes of heavy quarks have been computed¹².

2 Top quark mass determination

For the top quark system the relation between the resonance energy and the quark mass is given by

$$E_{\text{res}} = 2m_t + E_1^{\text{p.t.}} + \delta^{\Gamma_t} E_{\text{res}}, \quad (1)$$

where $E_1^{\text{p.t.}}$ is the perturbative contribution to the ground state energy and $\delta^{\Gamma_t} E_{\text{res}} = 100 \pm 10$ MeV is the effect of higher-order resonances and the finite width of the top quark. Non-perturbative effects are negligible.

To evaluate the perturbative contribution we use the result of Ref.⁷ for $E_1^{\text{p.t.}}$ up to N³LO. In Fig. 1 $E_1^{\text{p.t.}}$ is plotted in NLO, NNLO and N³LO approximation as a function of the renormalization scale of the strong coupling

^aNote that this corresponds to the N³LL corrections to the spin-dependent part of the energy levels.




Figure 1: Ground state energy $E_1^{\text{p.t.}}$ of the $t\bar{t}$ bound state in the zero-width approximation as a function of the renormalization scale μ for $S = 1$. The short-dashed (blue), long-dashed (green) and solid (red) line corresponds to the NLO, NNLO and N^3LO approximations.

constant for $S = 1$. One can see, that the N^3LO result shows a much weaker dependence on μ than the NNLO one. Moreover at the scale $\mu \approx 15$ GeV, which is close to the physically motivated soft scale $\mu_s \approx 30$ GeV, the N^3LO correction vanishes and furthermore becomes independent of μ , *i.e.* the N^3LO curve shows a local minimum. This suggests the convergence of the series for the ground state energy in the pole mass scheme. Similar results for the $\overline{\text{MS}}$ quark mass can be found in Ref. ¹³.

In Ref. ⁷ a universal relation has been obtained between the resonance energy in $t\bar{t}$ threshold production in e^+e^- annihilation or $\gamma\gamma$ collisions and the top quark pole mass

$$E_{\text{res}} = (1.9833 + 0.007 \frac{m_t - 174.3 \text{ GeV}}{174.3 \text{ GeV}} \pm 0.0009) \times m_t. \quad (2)$$

The central value is computed for $m_t = 174.3$ GeV. For the estimate of the theoretical uncertainty in Eq. (2) we assume a $\pm 100\%$ error in the Padé approximation of the still unknown three-loop static potential coefficient and vary the normalization scale in the stability interval which is roughly given by $0.4\mu_s < \mu < \mu_s$. Furthermore we use $\alpha_s(M_Z) = 0.1185 \pm 0.002$ and take into account the ± 10 MeV uncertainty in $\delta^{\Gamma_t} E_{\text{res}}$. Due to the very nice behaviour of the perturbative expansion for the ground state energy we do not expect large higher order corrections to our result. We want to mention that m_t extracted from Eq. (2) leads to a fixed-order quark mass which, by definition, does not suffer from renormalon ambiguities. Furthermore, it can be converted with high accuracy to the $\overline{\text{MS}}$ quark mass.




Figure 2: HFS of 1S bottomonium (a) and charmonium (b) as a function of the renormalization scale μ in the LO (dotted line), NLO (dashed line), LL (dot-dashed line), and NLL (solid line) approximations. For the NLL result, the (yellow) band reflects the errors due to $\alpha_s(M_Z) = 0.118 \pm 0.003$. The horizontal (green) band in (b) gives the experimental value 117.7 ± 1.3 MeV.

3 Prediction of $M(\eta_b)$

A compact analytical expression for the HFS to NLL, $E_{\text{hfs}}^{\text{NLL}}$, is given in Eq. (1) of Ref. ¹⁰. $E_{\text{hfs}}^{\text{NLL}}$ is a function of $y = \alpha_s(\mu)/\alpha_s(m_b)$.

In Fig. 2(a), the HFS for the bottomonium ground state is plotted as a function of μ in the LO, NLO, LL, and NLL approximations. As can be seen, the LL curve shows a weaker scale dependence compared to the LO one. The scale dependence of the NLO and NLL expressions is further reduced, and, moreover, the NLL approximation remains stable up to smaller scales than the fixed-order calculation. At the scale $\mu' \approx 1.3$ GeV, which is close to the inverse Bohr radius, the NLL correction vanishes. Furthermore, at $\mu'' \approx 1.5$ GeV, the result becomes independent of μ ; *i.e.*, the NLL curve shows a local maximum. This suggests a nice convergence of the logarithmic expansion despite the presence of the ultrasoft contribution with α_s normalized at the rather low scale $\bar{\mu}^2/m_b \sim 0.8$ GeV. By taking the difference of the NLL and LL results at the local maxima as a conservative estimate of the error due to uncalculated higher-order contributions, we get $E_{\text{hfs}} = 39 \pm 8$ MeV. A similar error estimate is obtained by the variation of the normalization scale in the physically motivated soft region 1 – 3 GeV.

So far only the perturbative part of the HFS has been discussed. In Ref. ¹⁰ it has been argued that the non-perturbative contributions are small. On one hand this is supported by the good agreement of our result with recent lattice analysis (see discussion in Ref. ¹⁰). On the other hand, in the charm system very good agreement of the HFS with the experimental data is obtained as can be seen in Fig. 2(b) where the experimental value 117.7 ± 1.3 MeV is very close to the theoretical value at the local maximum of the NLL curve which reads

$E_{\text{hfs}} = 104$ MeV. We should emphasize the crucial role of the resummation to bring the perturbative prediction closer to the experimental figure.

Our final prediction of the mass of the as-yet undiscovered η_b meson reads

$$M(\eta_b) = 9421 \pm 11 \text{ (th)} \pm_8^+ (\delta\alpha_s) \text{ MeV} , \quad (3)$$

where the errors due to the high-order perturbative corrections and the non-perturbative effects are added up in quadrature in “th”, whereas “ $\delta\alpha_s$ ” stands for the uncertainty in $\alpha_s(M_Z) = 0.118 \pm 0.003$. If the experimental error in future measurements of $M(\eta_b)$ will not exceed a few MeV, the bottomonium HFS will become a competitive source of $\alpha_s(M_Z)$ with an estimated accuracy of ± 0.003 , as can be seen from Fig. 2.

Acknowledgments

I would like to thank B. Kniehl, A. Penin, A. Pineda and V. Smirnov for a very pleasant and fruitful collaboration. This work was supported by HGF Grant No. VH-NG-008.

References

1. M. Martinez and R. Miquel, Eur. Phys. J. C **27** (2003) 49.
2. A.H. Hoang et al., Eur. Phys. J. direct C **3**, 1 (2000).
3. A.H. Hoang, A.V. Manohar, I.W. Stewart, and T. Teubner, Phys. Rev. Lett. **86** (2001) 1951.
4. A.H. Hoang, Phys. Rev. D **69** (2004) 034009.
5. A. Pineda and J. Soto, Nucl. Phys. Proc. Suppl. **64** (1998) 428.
6. B.A. Kniehl, A.A. Penin, V.A. Smirnov, and M. Steinhauser, Nucl. Phys. B **635** (2002) 357.
7. A.A. Penin and M. Steinhauser, Phys. Lett. B **538** (2002) 335.
8. B.A. Kniehl and A.A. Penin, Nucl. Phys. **B577**, 197 (2000); A.V. Manohar and I.W. Stewart, Phys. Rev. D **63**, 054004 (2001).
9. B.A. Kniehl, A.A. Penin, V.A. Smirnov, and M. Steinhauser, Phys. Rev. Lett. **90** (2003) 212001; Erratum *ibid.* **91** (2003) 139903.
10. B.A. Kniehl, A.A. Penin, A. Pineda, V.A. Smirnov, and M. Steinhauser, Phys. Rev. Lett. **92** (2004) 242001.
11. A.A. Penin, A. Pineda, V.A. Smirnov, and M. Steinhauser, hep-ph/0403080, to appear in Phys. Lett. B.
12. A.A. Penin, A. Pineda, V.A. Smirnov, and M. Steinhauser, hep-ph/0406175, to appear in Nucl. Phys. B.
13. Y. Kiyo and Y. Sumino, Phys. Rev. D **67** (2003) 071501.